NOBANIS - Marine invasive species in Nordic waters - Fact Sheet

Monocorophium sextonae

Author of this species fact sheet: Kathe R. Jensen, Zoological Museum, Natural History Museum of Denmark, Universiteteparken 15, 2100 København Ø, Denmark. Phone: +45 353-21083, E-mail: <u>krjensen@snm.ku.dk</u>

Bibliographical reference – how to cite this fact sheet:

Jensen, Kathe R. (2015): NOBANIS – Invasive Alien Species Fact Sheet – *Monocorophium sextonae* – From: Identification key to marine invasive species in Nordic waters – NOBANIS www.nobanis.org, Date of access x/x/201x.

Species description

Species name

Monocorophium sextonae, (Crawford, 1937) - a mudshrimp

Synonyms

Corophium sextonae. According to World Register of Marine Species (WORMS), the accepted name is *Monocorophium sextonae*.

Common names

Mudshrimp (corophiids) (UK); Slijkgarnaal (NL); Sextonkrebs (NO).

Identification

This species is only 5 mm long and therefore identification needs microscopic examination. Urosome segments are fused and uropod 1 with lateral spines but no setae. \Im antenna 2 with long spine distally on peduncle.

Distribution

Native area

New Zealand has been claimed to be the native area because the oldest confirmed specimens are from this area (Hurley, 1954). It has sometimes been claimed to be a native of the NE Atlantic and invasive in New Zealand (Costello, 1993). The earliest record from New Zealand is from 1921 (Wolff, 2005), and hence this may not be the native area either. Thus it should probably be called cryptogenic (Kerckhof et al., 2007).

Introduced area

Monocorophium sextonae was introduced to the UK near Plymouth in the 1930s, where it was described as a new species. It had not been present in that area in previous samples (1895-1911), which is why it was considered introduced already at the time of its description (Hurley, 1954). It was introduced to Ireland in the 1970s (Costello, 1993), and also to Scotland (Moore, 1980). According to Wolff (2005) there is a record from Portugal in 1930. It seems to be common in the Mediterranean (Ponti et al., 2002; Magni et al., 2004; Guerra-Garcia & Garcia-Gomez, 2009). It is

established on the coast of Belgium, where it is associated with shipwrecks (Kerckhof, 2007), on the Atlantic coast of France (Goulletquer et al., 2002), and in the Netherlands, the first record being from 1952 (Platvoet & Pinkster, 1995; Wolff, 2005). In Germany it has been found at the island Sylt in the Wadden Sea (Nehring & Leuchs, 1999), but it is uncertain whether it is established. It also occurs on the Skagerrak coast of Norway, where it was first reported in 1985 (Brattegard & Holthe, 1997).

Furthermore, it occurs in Tasmania, Australia, though this may be part of its native area.

Vector

Probably hull fouling (Hurley, 1954), but transfer with oysters is another possibility (Nehring & Leuchs, 1999).

Ecology

Habitat

This species is often associated with man-made structures, such as oil platforms or harbours (Ponti et al., 2002; Guerra-Garcia & Garcia-Gomez, 2009). It builds mud-tubes, which are attached to seaweeds, fouling organisms such as sponges and ascidians, or artificial substrates.

Reproduction

Females brood the eggs. The only record mentions 33 eggs in a brood (Hurley, 1954).

Impacts

None described. Interspecific competition seems likely; a native species of *Corophium* decreased in abundance after the introduction of *M. sextonae* in Ireland (Costello, 1993).

References

- Bachmann, V., Cegielka, E., Wagner, P., Usseglio-Polatera, P. and Moreteau, J.-C. 1995. Installation de l'amphipode *Corophium curvispinum* et de la palourde asiatique *Corbicula* sp. dans partie française de la Meselle. Hydroécologie Appliquée 7(1-2): 185-190.
- Brattegard, T. and Holthe, T. 1997. Distribution of marine, benthic macro-organisms in Norway. Utredning for Direktoratet for naturforvaltning, Trondheim, 380 pp.
- Costello, M.J. 1993. Biogeography of alien amphipods occurring in Ireland, and interactions with native species. Proceedings of the First European Crustacean Conference, 1992. Crustaceana 65(3): 287-299.
- Främmande arter 2006. *Corophium curvispinum*. Fact sheet. Available at: <u>http://www.frammandearter.se/0/2english/pdf/Corophium curvispinum.pdf</u> (last accessed on 26 August 2009).

Främmande arter 2006. *Monocorophium sextonae*. Fact sheet. Available at: <u>http://www.frammandearter.se/0/2english/pdf/Corophium_sextonae.pdf</u> (last accessed on 26 August 2009).

- Goulletquer, P., Bachelet, G., Sauriau, P.G. and Noel, P. 2002. Open Atlantic coast of Europe a century of introduced species into French waters. In: Invasive aquatic species of Europe. Distribution, impacts and management (eds. E. Leppäkoski, S. Gollasch and S. Olenin), pp. 276-290. Kluwer Academic Publishers, Dordrecht, The Netherlands.
- Guerra-Garcia, J.M. and Garcia-Gómez, J.C. 2009. Recolonization of macrofauna in unpolluted sands in a polluted harbour : A field approach using experimental trays. Estuarine, Coastal and Shelf Science 81: 49-58.
- Hartog, C. den, van den Brink, F.W.B. and vander Velde, G. 1992. Why was the invasion of the river Rhine by *Corophium curvispinum* and *Corbicula* species so successful? Journal of Natural History 26(6): 1121-1129.
- Herkül. K. and Kotta, J. 2007. New records of the amphipods Chelicorophium curvispinum, Gammarus tigrinus, G. duebeni, and G. lacustris in the Estonian coastal sea. Proceedings of the Estonian Academy of Sciences, Biology and Ecology 56(4): 290-296.
- Hurley, D.E. 1954. Studies on the New Zealand amphipodan fauna. No. 7. The family Corophiidae, including a new species of *Paracorophium*. Transactions of the Royan Society of New Zealand 82(2): 431-460.
- Jażdżewski, K. and Konopacka, A. 2000. Immigration history and present distribution of alien crustaceans in Polish waters. In: The biodiversity crisis and Crustacea (eds. I.C. von Vaupel Klein and F.R. Schram). Crustacean Issues 12: 55-64.
- Kerckhof, F., Haelters, J. and Golasch, S. 2007. Alien species in the marine and brackisch ecosystem: the situation in Belgian waters. Aquatic Invasions 2(3): 243-257.
- Kotta, J., Herkül. K., Kotta, I. and Orav-Kotta, H. 2006. Invasion history and distribution of the key benthic alien invertebrate species in the Estonian coastal sea. In: Alien invasive species in the north-eastern Baltic Sea: population dynamics and ecological impacts (eds. H. Ojaveer and J. Kotta), pp. 12-17. Estonian Marine Institute Report Series No. 14, Tallinn.
- Magni, P., Micheletti, S., Casu, D., Floris, A., De Falco, G. and Castelli, A. 2004. Macrofaunal community structure and distribution in a muddy coastal lagoon. Chemistry and Ecology 20(S1): 397-409.
- Moon, H.P. 1970. Corophium curvispinum (Amphipoda) recorded again in the British Isles. Nature 226: 976.
- Moore, P.G. 1980. *Corophium sextonae* in Scottish waters. Journal of the Marine Biological Association of the United Kingdom 60: 1075.
- Nehring, S. and Leuchs, H. 1999. Neozoa (Makrozoobenthos) an der deutschen Nordseeküste Eine Übersicht. Bundesanstalt für Gewässerkunde, Koblenz, Bericht BfG-1200, 131pp.
- Platvoet, D. and Pinkster, S. 1995. Changes in the amphipod fauna (Crustacea) of the Rhine, Meuse and Scheldt estuary due to the 'Delta Plan' coastal engineering works. Netherlands Journal of Aquatic Ecology 29(1): 5-30.
- Ponti, M., Abbiati, M. and Ceccherelli, V.U. 2002. Drilling platforms as artificial reefs: distribution of macrobenthis assemblages of the "Paguro" wreck (northern Adriatic Sea). ICES Journal of Marine Science 59: S316-S323.
- Taylor, P.M. and Harris, R.R. 1986. Osmoregulation in *Corophium curvispinum* (Crustacea: Amphipoda), a recent coloniser of freshwater. I. Sodium ion regulation. Journal of Comparative Physiology B: Biochemical, Systemic, and Environmental Physiology 156(3): 323-329.
- Van den Brink, F.W.B., van der Velde, G. and bij de Vaate, A. 1993. Ecological aspects, explosive range extension and impact of a mass invader, *Corophium curvispinum* Sars, 1895 (Crustacea: Amphipoda), in the Lower Rhine (The Netherlands). Oecologia 93: 224-232.

Wolff, W.J. 2005. Non-indigenous marine and estuarine species in The Netherlands. Zoologische Mededelingen 79(1): 1-116.